

Diagnosing Chilling and Flooding Injury to Corn

Ross Ennen, Sr. Research Associate, and Mark Jeschke, Ph.D., Agronomy Manager

SUMMARY

- Farmers often plant corn very early to increase yield potential and to avoid delays later in the planting season.
- Early planting offers potential advantages, but it also carries greater risk of cold injury and damage from pests.
- Ultra-early planted corn may require up to four weeks to emerge, depending on soil and weather conditions.
- During this time, the seed and emerging seedling are highly vulnerable to damage from insects, diseases, and herbicide exposure. The emerging seedling may also encounter adverse field conditions such as crusting or ponding.
- In addition, chilling temperatures caused by rain, melting snow, or cold soils can damage the seed during imbibition or injure the delicate structures of the emerging seedling.
- These stresses are often compounded under no-till conditions due to lower soil temperatures and additional water in the crop residue.

INTRODUCTION

Choosing corn planting date is an important management practice for maximizing corn yield potential. Often that date is dictated by prevailing weather and soil conditions, as well as the size of acreage to be planted. Historically, corn planting dates have moved earlier due to lengthening of the frost-free season, improved stress tolerance of newer corn hybrids, and the desire to avoid planting delays that could reduce yields. Early-planted corn is subject to greater risk of encountering cold temperatures and adverse weather systems often associated with those early spring dates.

Corn is a warm-season crop with tropical origins. It is not surprising then, that corn is susceptible to stresses that result from early planting under cold soil conditions. When corn is planted extremely early and soil temperatures are below 50 °F (10 °C), it is possible for corn seeds to remain in the soil up to three to four weeks prior to emergence. The length of this period will depend on the soil

temperature and its water holding properties. During this time, corn may encounter a range of stresses, including injury from pre-emergence residual herbicides, insect damage, and disease pressure.



Snow covering a recently-planted corn field on May 1, 2013.

Even more problems may result from the physical properties of the seedbed, including crusting, ponding, or saturated soils. In addition, cold temperatures resulting from cold rain or even snow can severely impact the seed. This *Crop Bulletin* will discuss effects of cold soils and water on germination and emergence of corn, including diagnosing plant injury symptoms caused by chilling and flooding.

EFFECT OF COLD SOILS AND WATER

The early spring seedbed is a very unfavorable environment for corn seeds. Though dry seeds can be stored unharmed for many years at -20 °F (-29 °C) or below, corn planted very early is at risk to cold injury and even death once the seeds begin to imbibe water. Early planting often exposes seeds to hydration with cold water, which can cause direct physiological damage. In addition, prolonged exposure to low temperatures reduces seed and plant metabolism and vigor, increases sensitivity to herbicides and seedling blights and causes oxidation damage due to the effects of free radicals in the cell. Free radicals are unstable molecules that damage cells and organs. This damage is similar to

damage that occurs in mammalian cells during aging and sun exposure.

When the dry seed imbibes cold water as a result of a cold rain or melting snow, imbibitional chilling injury may result. The cell membranes of the seed lack fluidity at low temperatures, and under these conditions, the hydration process can result in rupture of the membranes. Cell contents then leak through this rupture and provide a food source for invading pathogens. Cold water can similarly affect seedling structures as they begin to emerge.

Corteva Agriscience routinely conducts research studies on corn germination and emergence in stressful environments, in fields where soil temperatures are at or below the minimum recommended threshold for planting corn. Results of these studies have shown that temperatures at or below 50 °F (10 °C) are often detrimental to the germination and emergence process, especially if they persist long after planting (Table 1).

Table 1. Planting dates, average soil temperature the week after planting, cumulative precipitation the week after planting, days to emergence, and final stand in Corteva Agriscience research plots in 2018.

Location	Plant Date	Soil Temp °F	Precip. in	Days to VE*	Stand %
Riverdale, MI**	April 11	38	1.19	23	73
Janesville, WI**	April 11	39	1.29	21	79
Johnston, IA**	April 12	42	1.54	19	83
Eau Claire, WI	April 27	51	0.46	10	89
Moorhead, MN	April 30	50	0.82	12	91
Olivia, MN	May 5	57	0.19	12	94
Flandreau, SD	May 4	NA	0.82	13	95

* Values reflect averages of multiple hybrids planted at each location
 ** Locations characterized as high stress environments for germination and emergence.

In Corteva research studies conducted in 2018, days to emergence and percent final stand varied considerably depending on the average soil temperature and rainfall during the week following planting. Three research locations experienced average soil temperatures below 50 °F (10 °C) with greater than 1 inch (2.5 cm) of rainfall the week following planting. These locations had substantially longer time to emergence and lower stand establishment than locations with soil temperatures above 50 °F (10 °C) and less rainfall. These data show that cold, wet soils after planting can have serious consequences for stand establishment. However, the degree of damage will vary

with soil type and is generally greater in heavier or poorly drained soils.

FLOODING EFFECTS ON EMERGENCE

Flooding can have an equally devastating effect on seedling emergence and survival as cold soils. Most corn hybrids can only survive for 24 to 48 hours under water, with smaller seedlings suffering the most damage. Flooding damages corn biochemically. By impairing mitochondria, it causes release of free radicals which damage cell membranes. Flooding also causes oxygen starvation and shifts the plant's metabolic processes to anaerobic fermentation. Resulting acidosis (low pH) can kill the cells. At a minimum, flooding reduces the plant's metabolic rate, making seedlings more sensitive to disease, insects, and herbicides. In fact, many disease-causing fungi such as *Pythium* thrive in standing water. Seedlings that are weakened by flooding or cold damage usually succumb to disease if the pathogen is present in the soil.

Flooding damage does not only occur in obvious ponded areas of a field. If fields are completely saturated to the soil surface and remain that way due to continual rain or limited drainage, seeds and non-emerged seedlings are under water. Flooding damage may occur in these areas just as in ponded areas.



Field with saturated soil following spring rainfall.

DIAGNOSING POOR STAND ESTABLISHMENT

Careful examination of damaged seedlings can provide clues into the likely causes of stand establishment problems following early planting or abnormally cold weather conditions. Table 2 lists the main symptoms and likely causes of early season damage. Figures 1-5 show diagnostic images of chilling and flooding damage to corn seedlings during germination and emergence.

CORTEVA AGRISCIENCE RESEARCH

For decades, plant breeders have selected within the natural variation expressed by corn genotypes to develop

hybrids with strong emergence and vigor characteristics under cool soil conditions. In the late 2000's, a new rating was introduced for corn products called stress emergence. Stress emergence is a measure of the genetic ability or potential to emerge in the stressful environmental conditions of cold, wet soils or short periods of severe low temperatures, relative to other products. Ratings of 7 to 9 indicate very good potential to establish normal stands under such conditions; a rating of 5 or 6 indicates average potential to establish normal stands under moderate stress conditions; and ratings of 1 to 4 indicate the product has below-average potential to establish normal stands under stress and should not be used if severe cold conditions are expected immediately after planting. Stress emergence is not a rating for seedling disease susceptibility, early growth, or speed of emergence.

Corteva research scientists are continuing to work to improve early season corn performance through conventional and molecular breeding, as well as through rigorous testing of research and commercial hybrids. By identifying molecular markers and pathways associated with superior cold germination, Corteva researchers are beginning to develop an understanding of the genetic basis of stress emergence. This knowledge should eventually lead to even stronger early-season performance in elite corn products.

Table 2. Corn seedling symptoms and likely causes.

Symptom	Likely Cause	Result
Stubby coleoptiles Leaves emerging prematurely	Imbibitional chilling or cold damage	Death, unless unprotected leaf reaches the surface
Brown tissue behind root tip Adventitious roots	Chilling damage Flooding	Chance for survival unless shoot meristem is damaged
Leafing underground Leaves growing along soil crust	Mechanical damage Soil crusting	Usually death, as seedlings lose ability to penetrate soil
Corkscrew mesocotyl or coleoptile	Temperature fluctuations Herbicide injury	Seedling death
Fused coleoptile or bursting on side	Cold damage Genetic tendency	Seedling death
Rotted seed or mesocotyl Spotty wilting	Seedling disease	Seedling death or stunting
Bleached leaves	Herbicide or cold injury	Seedlings can grow out of it unless impairment of

photosynthesis is extensive

Pruned roots	Insect damage	Weak seedlings, wilting
--------------	---------------	-------------------------



Figure 1. Imbibitional chilling and cold injury. Note club-shaped coleoptile and leafing out under-ground.



Figure 2. Corkscrew mesocotyl growth. Can be caused by cold soils, extreme soil temperature fluctuations, or soil crusting.



Figure 3. Fused coleoptile / bursting on the side caused by cold injury.



Figure 4.
Corn seedlings with necrotic tissue resulting from flooding.



Figure 5.
Corn seedlings showing both brown root tissue and bursting on the side due to cold and flooded soil conditions.

The foregoing is provided for informational use only. Please contact your Seed sales professional for information and suggestions specific to your operation. Product performance is variable and depends on many factors such as moisture and heat stress, soil type, management practices and environmental stress as well as disease and pest pressures. Individual results may vary. Our products are provided subject to the terms and conditions of purchase which are part of the labeling and purchase documents.

March 2020